

## ARTIGO / ARTÍCULO / ARTICLE

### *Trinoparvus majeri* sp. nov. from Thailand (Coleoptera: Dermestidae: Trinodinae: Trinoparvini).

Jiří Háva

Department of Forest Protection and Entomology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences  
Kamýcká 1176, CZ-165 21, Prague 6 - Suchdol, Czech Republic. e-mail: jh.dermestidae@volny.cz

**Abstract:** A third species of the genus *Trinoparvus* Háva, 2004, *Trinoparvus majeri* sp. nov. from Thailand is described, illustrated and compared with the two previously known species. The new species differs from *Trinoparvus laboriosus* Háva, 2004 and *T. villosus* Háva, 2004 by its bicolorous elytra.

**Key words:** Coleoptera, Dermestidae, Taxonomy, new species, Trinoparvini, *Trinoparvus*, Thailand.

**Resumen:** *Trinoparvus majeri* sp. nov. de Tailandia (Coleoptera: Dermestidae: Trinodinae: Trinoparvini). Se describe una tercera especie del género *Trinoparvus* Háva, 2004, *Trinoparvus majeri* sp. nov., de Tailandia, y se ilustra y compara con las dos especies previamente conocidas. La nueva especie se diferencia de *Trinoparvus laboriosus* Háva, 2004 y *T. villosus* Háva, 2004 por sus élitros bicoloros.

**Palabras clave:** Coleoptera, Dermestidae, Taxonomía, nueva especie, Trinoparvini, *Trinoparvus*, Tailandia.

Recibido: 12 de noviembre de 2014

Aceptado: 18 de noviembre de 2014

Publicado on-line: 29 de noviembre de 2014

urn:lsid:zoobank.org:pub:CCA48067-2A67-4555-B7A7-741438D3227D

## Introduction

The genus *Trinoparvus* was described by Háva (2004) to include two species from Madagascar and New Caledonia, respectively. The genus belongs to the subfamily Trinodinae, tribe Trinoparvini (Háva in Kirejtshuk et al. 2010, Háva 2014). In the present article, a third species recently collected in Thailand is described.

## Material and methods

The following measurements were made:

total length (TL) - linear distance from anterior margin of head to apex of elytra

pronotal length (PL) - maximum length measured from anterior margin to posterior margin

pronotal width (PW) - maximum linear transverse distance

elytral length (EL) - linear distance from shoulder to apex of elytron

elytral width (EW) - maximum linear transverse distance

All photos were made using a camera Olympus SZX16.

## Taxonomy

### Subfamily Trinodinae

#### Tribe Trinoparvini

##### *Trinoparvus majeri* sp. nov. (Figs. 1-5)

**Type locality.** Thailand, Ranong prov., Ban Na env.

**Type material.** Holotype (♂): "Thailand, 22-26.III.1996, Ranong prov., 9°34' 98°42', Ban Na env., K. Majer lgt.". Holotype specimen provided with label: "HOLOTYPE *Trinoparvus majeri* sp. nov. Jiří Háva det. 2014", and deposited in the National Museum, Prague (Czech Republic).

#### Description.

**Male.** Body (Figs. 1-3) brownish-black on dorsal and ventral surfaces, shiny, oval, with erect long light setation. Measurements (mm): TL 1.1 PL 0.2 PW 0.5 EL 0.7 EW 0.7. Head finely punctate with erect, long, light brown setation; maxillary palpi brown; pubescence on mentum denser. Ocellus on front present. Antennae yellow, 10-segmented, antennal club 3-segmented, segment XI about as long as broad, IX small and X obviously transverse (Fig. 4). Pronotum finely punctate as head, with erect, long, light brown setation, anterior angles not visible from above. Scutellum small, triangular without setation. Elytra finely punctate as head and pronotum with erect, long, light brown setation. Each elytron black, with two reddish circular spots. Epipleura short, black. Legs brown with long yellow setae. Prosternal process quite straight, broad, without tip. Metasternum with erect, long, light brown setation and without small striae near coxa. Abdominal sternites brown, finely punctate with long, light brown setation; first visible abdominal sternite with small striae near post-coxal line. Male genitalia (Fig. 5), length 0.35 mm.

**Female.** Unknown.

**Differential diagnosis.** The new species differs from the two known species *Trinoparvus laboriosus* Háva, 2004 (Madagascar) and *T. villosus* Háva, 2004 (New Caledonia) by its bicolorous elytra.

**Name derivation.** Dedicated in memoriam to my very good friend and collector of the new species †Karel Majer (1949-2000).

#### List of species of the genus *Trinoparvus* Háva, 2004

*Trinoparvus laboriosus* Háva, 2004: Madagascar

*Trinoparvus majeri* sp. nov.: Thailand

*Trinoparvus villosus* Háva, 2004: New Caledonia

## Acknowledgements

I am obliged to Jiří Hájek (National Museum, Prague) for loaning me the interesting material. The research was supported by the Internal Grant Agency (IGA n.20124364) Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague.


## References

---

HÁVA, J. 2004. World keys to the genera and subgenera of Dermestidae (Coleoptera), with descriptions, nomenclature and distributional records. *Acta Musei Nationalis Pragae, Series B, Natural History* 60: 149-164.

HÁVA, J. 2014. Dermestidae, Derodontidae, Jacobsoniidae, Nosodendridae. In: ZAHRADNÍK, P. & HÁVA, J. Catalogue of the world genera and subgenera of the superfamilies Derodontoidea and Bostrichoidea (Coleoptera: Derodontiformia, Bostrichiformia). *Zootaxa* 3754: 301-352.

KIREJTSCHUK, A.G.; HÁVA, J. & NEL, A. 2010. New genus and species of subfamily Trinodinae (Coleoptera, Polyphaga, Dermestidae) from Lowermost Eocene French Amber. *Zoosystematica Rossica* 19: 54-69.



Figs. 1-5.- *Trinoparvus majeri* sp. nov. (holotype).  
1.- Dorsal aspect. 2.- Frontal aspect. 3.- Lateral aspect. 4.- Antenna. 5.- Aedeagus.