

NOTA / NOTE

Stenolemus novaki Horváth, 1888 (Hemiptera, Reduviidae) in Portugal.

José Manuel Grosso-Silva¹ & Sónia Ferreira²

¹ Museu de História Natural e da Ciência da Universidade do Porto (MHNC-UP) / PRISC, Praça Gomes Teixeira, 4099-002 Porto, Portugal. e-mail: jmgrossosilva@mhnc.up.pt

² CIBIO-InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Vairão, Portugal.

Abstract: *Stenolemus novaki* Horváth, 1888 (Hemiptera, Reduviidae) is recorded from mainland Portugal for the second time, following a recent photographic record. A map synthesizing the knowledge about the species distribution in the Iberian Peninsula is included.

Key words: Hemiptera, Reduviidae, Emesinae, *Stenolemus novaki*, Iberian Peninsula.

Resumen: *Stenolemus novaki* Horváth, 1888 (Hemiptera, Reduviidae) en Portugal. Se cita *Stenolemus novaki* Horváth, 1888 (Hemiptera, Reduviidae) de Portugal continental por segunda vez, siguiendo un registro fotográfico reciente. Se incluye un mapa que sintetiza el conocimiento sobre la distribución de la especie en la Península Ibérica.

Palabras clave: Hemiptera, Reduviidae, Emesinae, *Stenolemus novaki*, Península Ibérica.

Recibido: 21 de agosto de 2020

Publicado on-line: 27 de septiembre de 2020

Aceptado: 17 de septiembre de 2020

Stenolemus novaki Horváth, 1888 (Hemiptera, Reduviidae), the only representative of the genus *Stenolemus* Signoret, 1858 in Europe, is an emesine assassin bug described from Croatia ("Dalmatia") and Italy (Lesina) by HORVÁTH (1888).

According to PUTSHKOV & MOULET (2009), who conducted a thorough compilation of the available ecological data, *Stenolemus novaki* is probably a xerothermic and heliophilous species with a preference for rocky areas.

Up to the present, the known European distribution of the species includes Bulgaria, Croatia, France, Greece, Italy, Macedonia, and Spain, while outside Europe it has been recorded from the Canary Islands, Egypt, Israel, Saudi Arabia, Sudan, and Turkey (WYGODZINSKY, 1966; PUTSHKOV & PUTSHKOV, 1996; PUTSHKOV & MOULET, 2009; SIMOV et al., 2017).

The current known Iberian distribution includes the Spanish provinces of Barcelona (RIBES, 1961 and several additional records, cf. PUTSHKOV & MOULET, 2009) and Girona, specifically from the Medes Islands, from where ESPAÑOL (1984) tentatively identified several nymphs. In Portugal, the only available information pertains to a very recent record from the district of Setúbal [specific location: Setúbal (municipality): Castanhos (38.5231° N, -8.9997° W; 29SNC0063), Parque Natural da Arrábida], based on pictures taken by Pedro Silva on 13/11/2019 and housed in the web platform Biodiversity4All (<https://www.biodiversity4all.org/observations/35646460>) and GBIF (UEDA, 2020).

In this contribution we record *Stenolemus novaki* from south-western Portugal, collected within the scope of the InBIO Barcoding Initiative, confirming its presence and extending southwards its known distribution in the country as shown in Map 1, which marks the Portuguese districts where the species was recorded and summarizes its Spanish (province level) distribution just based on literature

compilation. In the material examined section, as in the above-mentioned record from Setúbal, the dates presented are in the Day/Month/Year format and the coordinates are in the Military Grid Reference System (MGRS).

Material examined

PORUGAL: Beja district: Odemira municipality: Vale de Gomes, 1,7 km WSW of Algoceira (37,6116° N, 8,7134° W; 29SNB2562, 4 m above sea level), Parque Natural do Sudoeste Alentejano e Costa Vicentina, 27/05/2019, 1 ♂, Sónia Ferreira leg., InBIO Barcoding Initiative reference collection (INV08667) (Vairão, Portugal).

Discussion

The known distribution range of *Stenolemus novaki* Horváth, 1888 in the Iberian Peninsula was remarkably extended with the aforementioned record obtained by Pedro Silva and is further increased with the new record presented herein (Map 1).

As in the case of *Lachnaia (Lachnaia) variolosa* (Linnaeus, 1767), recently recorded from Portugal for the first time with a significant westwards range expansion (GROSSO-SILVA et al., 2020), *Stenolemus novaki* is probably present in suitable areas in a number of Spanish provinces located between Barcelona and the Portuguese south-western coast, its small size and elusive behaviour possibly accounting for the lack of records in such an extensive area of the Iberian Peninsula.

Acknowledgments

InBIO Barcoding Initiative is funded by the European Union's Horizon 2020 Research and Innovation programme under grant agreement No 668981 and by the project PORBIOTA - Portuguese E-Infrastructure for Information and Research on Biodiversity (POCI-01-0145-FEDER-022127), supported by Operational Thematic Program for Competitiveness and Internationalization (POCI), under the PORTUGAL 2020 Partnership Agreement, through the European Regional Development Fund (FEDER).

References

- ESPAÑOL, F. 1984. *Els insectes de les Illes Medes*, pp. 231-244. In: ROS, J.; OLIVELLA, I. & GILI, J. M. (eds.). *Els Sistemes naturals de les illes Medes*. Arxius de la Secció de Ciències, LXXIII. Institut d'Estudis Catalans, Barcelona. 826 pp.
- GROSSO-SILVA, J.M.; SOARES, A.; MONTEIRO, E.; SANTOS, R.; FÉLIX, R.; ANTUNES, S. & GARCIA-PEREIRA, P. 2020. *Lachnaia (Lachnaia) variolosa* (Linnaeus, 1767) (Coleoptera, Chrysomelidae), new species for Portugal. *Arquivos Entomológicos*, 22: 9-10.
- HORVÁTH, G. 1888. Matériaux pour servir à l'étude des Hemiptères de la faune paléarctique. *Revue d'Entomologie*, VII: 168-189.
- PUTSHKOV, P.V. & MOULET, P. 2009. *Hémiptères Reduviidae d'Europe Occidentale*. Faune de France, 92. France et régions limitrophes. Fédération Française des Sociétés de Sciences Naturelles, Paris. 668 pp. + 24 planches.

PUTSHKOV, P.V. & PUTSHKOV, V.G. 1996. Family Reduviidae Latreille, 1807 - assassin-bugs, pp. 148-265. In: AUKEMA, B. & RIEGER, C. (eds.). Catalogue of the Heteroptera of the Palaearctic Region. Vol. 2. Cimicomorpha I. The Netherlands Entomological Society, Amsterdam. 361 pp.

RIBES, J. 1961. I. - Contribución al estudio de los Reduviidae de Cataluña. *Miscelánea Zoológica*, 1(4): 57-73.

SIMOV, N.; GRADINAROV, D. & DAVRANOGLOU, L.-R. 2017. Three new assassin bug records (Hemiptera: Heteroptera: Reduviidae) for the Balkan Peninsula. *Ecologica Montenegrina*, 13: 25-29.

UEDA, K. 2020. iNaturalist Research-grade Observations. iNaturalist.org. Occurrence dataset <https://doi.org/10.15468/ab3s5x> accessed via GBIF.org on 2020-08-19. <https://www.gbif.org/occurrence/2456578642>

WYGODZINSKY, P.W. 1966. A monograph of the Emesinae (Reduviidae, Hemiptera). *Bulletin of the American Museum of Natural History*, 133: 1-616.



Map 1.- Iberian distribution of *Stenolemus novaki* Horváth, 1888, with the previously known Portuguese districts and Spanish provinces (in grey), the new Portuguese districts (in blue), and the MGRS 10x10 km squares of the collecting sites in Portugal (new sites in orange, previous sites in green).